

The role of RCAN1 in Alzheimer pathogenesis

<https://neurodegenerationresearch.eu/survey/the-role-of-racan1-in-alzheimer-pathogenesis/>

Principal Investigators

Song, Weihong

Institution

University of British Columbia

Contact information of lead PI

Country

Canada

Title of project or programme

The role of RCAN1 in Alzheimer pathogenesis

Source of funding information

CIHR

Total sum awarded (Euro)

€ 594,750

Start date of award

01/07/2015

Total duration of award in years

5.0

The project/programme is most relevant to:

Alzheimer's disease and other dementias

Keywords

alzheimer's disease| amyloid beta protein| diabetes| drug dedevelopment| high glucose| neuronal apoptosis| racn1| transgeneic mice

Research Abstract

Lay Summary

Alzheimer's disease (AD) is the most common neurodegenerative disorder leading to dementia in the elderly population. Patients with diabetes have a greater risk to develop AD and a majority of AD patients also have hyperglycemia. The mechanism linking diabetes to AD remains

undefined. Progressive neuronal loss is one of the pathological features of AD. People with Down Syndrome having extra copy of chromosome 21 develop characteristic AD neuropathology after middle age and several genes on chromosome 21 including APP and the regulator of calcineurin 1 (RCAN1) are implicated in AD pathogenesis. Our preliminary studies suggest that RCAN1 promotes neurodegeneration and our working hypothesis is that abnormal RCAN1 expression and function contribute to Alzheimer pathogenesis. This proposal seeks to elucidate the molecular mechanism of RCAN1's effect on Alzheimer pathogenesis and define its role in linking diabetes to AD. The results will be of clinical significance and provide evidences for targeting RCAN1 in AD treatment.

Further information available at:

Types:

N/A

Member States:

Canada

Diseases:

Alzheimer's disease & other dementias

Years:

2016

Database Categories:

N/A

Database Tags:

N/A